

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Chicken IgG anti-Goat IgG (H+L)-Biotin, MinX none, Gallus, Polyclonal DNA-SEC-182896

| | |
|----------------------------|---|
| Article Name | Chicken IgG anti-Goat IgG (H+L)-Biotin, MinX none, Gallus, Polyclonal |
| Biozol Catalog Number | DNA-SEC-182896 |
| Supplier Catalog Number | SEC-182896 |
| Alternative Catalog Number | DNA-SEC-182896 |
| Manufacturer | dianova |
| Host | Gallus |
| Category | Antikörper |
| Application | ELISA |
| Species Reactivity | Goat |
| Immunogen | Goat IgG whole molecule |
| Conjugation | Biotin |
| Product Description | Anti-Goat IgG Biotin Antibody generated in chicken detects goat IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well a... |
| Clonality | Polyclonal |
| Concentration | 1.0 mg/mL |
| Isotype | Ig |
| Buffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |

| | |
|--------------------|--|
| Purity | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Chicken Serum, Goat IgG and Goat Serum. |
| Formula | 20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3 |
| Target | Goat |
| Antibody Type | Polyclonal Antibody |
| Application Dilute | WB: 1:2,000 - 1:10,000 |
| Application Notes | Anti-Goat IgG Biotin Antibody has been tested by ELISA and is assayed against 1.0 ug of Goat IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin S000-03 and ABTS (2,2-azino-bis-[3-ethylbenthiiazoline-6-sulfonic acid]) code ABTS-10 |