

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat IgG anti-Guinea Pig IgG (Fc)-Alk. Phos., MinX none, ALP, Polyclonal , AP DNA-SEC-182912

| | |
|----------------------------|---|
| Article Name | Goat IgG anti-Guinea Pig IgG (Fc)-Alk. Phos., MinX none, ALP, Polyclonal , AP |
| Biozol Catalog Number | DNA-SEC-182912 |
| Supplier Catalog Number | SEC-182912 |
| Alternative Catalog Number | DNA-SEC-182912 |
| Manufacturer | dianova |
| Host | Goat |
| Category | Antikörper |
| Application | DOT, ELISA |
| Species Reactivity | Guinea pig |
| Immunogen | Guinea Pig IgG F(c) fragment |
| Conjugation | AP |
| Product Description | Anti-Guinea Pig IgG F(c) Alkaline Phosphatase generated in goat is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc po... |
| Clonality | Polyclonal |
| Concentration | 1.0 mg/mL |
| Isotype | Ig |

| | |
|--------------------|--|
| Buffer | 0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0 |
| Purity | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Guinea Pig IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a s |
| Formula | 50 mM TrisHCl,150 mM NaCl,1 mM MgCl,0,1 mM ZnCl,50% (v/v) Glycerol,pH 8,0,sterile filtered,0,01% NaN3 |
| Target | Guinea Pig |
| Antibody Type | Polyclonal Antibody |
| Application Dilute | ELISA Dilution: 1:2,000 - 1:10,000, Immunohistochemistry Dilution: 1:200 - 1:1,000, Western Blot Dilution: 1:500 - 1:2,500 |
| Application Notes | Anti-Guinea Pig IgG F(c) Alkaline Phosphatase has been tested by ELISA and dot blot. This product has been assayed against 1.0 ug of Guinea Pig IgG in a standard capture ELISA using pNPP p-nitrophenyl phosphate code NPP-10 as a substrate for 30 minutes |