

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat IgG anti-Horse IgG (Fc)-unconj., MinX none, Polyclonal, Unconjugated DNA-SEC-182958

| Article Name | Goat IgG anti-Horse IgG (Fc)-unconj., MinX none, Polyclonal , Unconjugated |
|----------------------------|---|
| Biozol Catalog Number | DNA-SEC-182958 |
| Supplier Catalog Number | SEC-182958 |
| Alternative Catalog Number | DNA-SEC-182958 |
| Manufacturer | dianova |
| Host | Goat |
| Category | Antikörper |
| Application | DOT, ELISA |
| Species Reactivity | Equine |
| Immunogen | Horse IgG F(c) fragment |
| Conjugation | Unconjugated |
| Product Description | Horse IgG F(c) antibody generated in goat recognizes the F(c) portion of horse IgG. It is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. R |
| Clonality | Polyclonal |
| Concentration | 2.3 mg/mL |
| Isotype | lg |

| Buffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
|--------------------|--|
| Purity | Anti-Horse IgG F(c) was prepared from monospecific antiserum by immunoaffinity chromatography using Horse IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Horse IgG, Horse IgG F(c) |
| Formula | 20 mM K3PO4,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3 |
| Target | Horse |
| Antibody Type | Polyclonal Antibody |
| Application Dilute | ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000 |
| Application Notes | Anti-Horse IgG F(c) antibody has been tested by ELISA and dot blot and is suitable for Western Blot and Immunohistochemistry applications. Antibody should be optimized by end user for specific reactive conditions. |