

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat IgG anti-Horse IgG (Fc)-Biotin, MinX none, Polyclonal DNA-SEC-182970

Article Name	Goat IgG anti-Horse IgG (Fc)-Biotin, MinX none, Polyclonal
Biozol Catalog Number	DNA-SEC-182970
Supplier Catalog Number	SEC-182970
Alternative Catalog Number	DNA-SEC-182970
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	WB, IHC, ELISA
Species Reactivity	Equine
Immunogen	Horse IgG F(c) fragment
Conjugation	Biotin
Product Description	Anti-Horse IgG F(c) generated in goat is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc portion of horse IgG and oft...
Clonality	Polyclonal
Concentration	2.0 mg/mL
Isotype	Ig
Buffer	0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Horse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single
Formula	10 mM NaPO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Thimerosal
Target	Horse
Antibody Type	Polyclonal Antibody
Application Dilute	ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000
Application Notes	This product has been assayed against 1.0 ug of Horse IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin S000-03 and ABTS (2,2-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code ABTS-100 as a substrate for 30 minutes at room