

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat IgG anti-Human IgG (H+L)-FITC, MinX Bo,Ck,Go,Gp,Hm,Ho,Ms,Rb,Rt,Sh, Polyclonal DNA-SEC-182991

| | |
|----------------------------|---|
| Article Name | Goat IgG anti-Human IgG (H+L)-FITC, MinX Bo,Ck,Go,Gp,Hm,Ho,Ms,Rb,Rt,Sh, Polyclonal |
| Biozol Catalog Number | DNA-SEC-182991 |
| Supplier Catalog Number | SEC-182991 |
| Alternative Catalog Number | DNA-SEC-182991 |
| Manufacturer | dianova |
| Host | Goat |
| Category | Antikörper |
| Application | DOT |
| Species Reactivity | Human |
| Immunogen | Human IgG whole molecule |
| Conjugation | FITC |
| Product Description | Anti-Human IgG (H&L) Fluorescein generated in goat detects human Immunoglobulin G (IgG), both heavy and light chains of the antibody molecule are present. It is a protein complex composed of four peptide chains - two identical heavy chains and two id... |
| Clonality | Polyclonal |
| Concentration | 1.0 mg/mL |
| Isotype | Ig |

| | |
|--------------------|--|
| Buffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| Purity | Secondary Antibodies are prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in |
| Formula | 20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3 |
| Target | Human |
| Antibody Type | Polyclonal Antibody |
| Application Dilute | FLISA Dilution: 1:10,000 - 1:50,000, Flow Cytometry Dilution: 1:500 - 1:2,500, Fluorochrome Protein Value: 2.19, IF Microscopy Dilution: 1:1,000 - 1:5,000 |
| Application Notes | Anti-Human IgG (H&L) Fluorescein Conjugated Antibody has been tested by dot blot. Secondary antibodies aid in the detection or sorting of target antigens by binding to the primary antibody. Anti-Human IgG (H&L) Fluorescein antibody is ideal for Flow Cyto |