

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Goat IgG anti-Human IgG (Fc)-unconj., MinX none, Polyclonal, Unconjugated DNA-SEC-183004

Article Name	Goat IgG anti-Human IgG (Fc)-unconj., MinX none, Polyclonal , Unconjugated
Biozol Catalog Number	DNA-SEC-183004
Supplier Catalog Number	SEC-183004
Alternative Catalog Number	DNA-SEC-183004
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	DOT, ELISA
Species Reactivity	Human
Immunogen	Human IgG F(c) fragment
Conjugation	Unconjugated
Product Description	Anti-Human IgG F(c) generated in goat detects Human F(c). A proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc portion of
Clonality	Polyclonal
Concentration	5.17 mg/mL
Isotype	lg

Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	Anti-Human IgG F(c) Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resu
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3
Target	Human
Antibody Type	Polyclonal Antibody
Application Dilute	ELISA Dilution: 1:20,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:10,000 - 1:50,000
Application Notes	Goat Anti-Human IgG F(c) Antibody has been tested by ELISA and dot blot and is suitable for western blot, immunohistochemistry, or immunoassays where specificity to the immunoglobulin Fc region is desired. Optimal concentrations in immunoassays should be