

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit IgG anti-Human IgG (H+L)-HRPO, MinX Ms, Polyclonal DNA-SEC-183075

| Article Name | Rabbit IgG anti-Human IgG (H+L)-HRPO, MinX Ms, Polyclonal |
|----------------------------|--|
| Biozol Catalog Number | DNA-SEC-183075 |
| Supplier Catalog Number | SEC-183075 |
| Alternative Catalog Number | DNA-SEC-183075 |
| Manufacturer | dianova |
| Host | Rabbit |
| Category | Antikörper |
| Application | ELISA |
| Species Reactivity | Human |
| Immunogen | Anti-Human IgG (H&L) was produced by repeated immunization with human IgG whole molecule in rabbit. |
| Conjugation | HRP |
| Product Description | Anti-Human IgG (H&L) Peroxidase generated in rabbit detects human Immunoglobulin G (IgG), both heavy and light chains of the antibody molecule are present. It is a protein complex composed of four peptide chains - two identical heavy chains and two i |
| Clonality | Polyclonal |
| Concentration | 2.0 mg/mL |
| Isotype | Ig |
| Buffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |

| Purity | Anti-Human IgG (H&L) peroxidase conjugated antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by imm |
|--------------------|--|
| Formula | 20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin |
| Target | Human |
| Antibody Type | Polyclonal Antibody |
| Application Dilute | ELISA Dilution: 1:50,000 - 1:100,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:1,000 - 1:5,000 |
| Application Notes | Anti-Human IgG HRP has been tested by ELISA and is suitable for immunoblotting, western blot, dot blot, ELISA, immunoperoxidase electron microscopy and immunohistochemistry as well as other peroxidase-antibody based enzymatic assays requiring lot-to-lot |