

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Goat IgG anti-Mouse IgG+IgM+IgA (H+L)-unconj., MinX none, Polyclonal , Unconjugated DNA-SEC-183121

|                            |   |
|----------------------------|---|
| Article Name               | Goat IgG anti-Mouse IgG+IgM+IgA (H+L)-unconj., MinX none, Polyclonal , Unconjugated   |
| Biozol Catalog Number      | DNA-SEC-183121  |
| Supplier Catalog Number    | SEC-183121  |
| Alternative Catalog Number | DNA-SEC-183121  |
| Manufacturer               | dianova   |
| Host                       | Goat  |
| Category                   | Antikörper  |
| Application                | ELISA   |
| Species Reactivity         | Mouse   |
| Immunogen                  | Mouse IgG IgA and IgM whole molecule  |
| Conjugation                | Unconjugated  |
| Product Description        | Anti-Mouse IgG IgA and IgM whole molecule antibody generated in goat detects specifically Mouse IgG IgA and IgM whole molecules. This secondary antibody anti-Mouse is ideal for investigators who routinely perform ELISA, Sandwich ELISA, titration assa... |
| Clonality                  | Polyclonal  |
| Concentration              | 2.20 mg/mL  |
| Isotype                    | Ig  |

|                    |  |
|--------------------|--|
| Buffer             | 0.125 M Sodium Borate, 0.075 M Sodium Chloride, 0.005 M EDTA, pH 8.0   |
| Purity             | Anti-Mouse IgG IgA IgM (H&L) Antibody was prepared from polyspecific antiserum by immunoaffinity chromatography using antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophore |
| Formula            | 125 mM Sodium Borate, 75 mM NaCl, 5 mM EDTA, pH 8.0, sterile filtered, 0.01% NaN <sub>3</sub>  |
| Target             | Mouse  |
| Antibody Type      | Polyclonal Antibody  |
| Application Dilute | ELISA Dilution: 1:20,000 - 1:50,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000   |
| Application Notes  | Anti-Mouse IgG IgA IgM (H&L) Antibody has been tested by ELISA and is suitable for immunoprecipitation, immunodiffusion, conjugation and most immunological methods requiring lot-to-lot consistency, high titer and specificity. Specific conditions for reac |