

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Goat IgG anti-Mouse IgG+IgM+IgA (H+L)-unconj., MinX Bo,Ho,Hu, Polyclonal, Unconjugated DNA-SEC-183122

| Article Name               | Goat IgG anti-Mouse IgG+IgM+IgA (H+L)-unconj., MinX Bo,Ho,Hu, Polyclonal , Unconjugated  |
|----------------------------|--|
| Biozol Catalog Number      | DNA-SEC-183122   |
| Supplier Catalog Number    | SEC-183122   |
| Alternative Catalog Number | DNA-SEC-183122   |
| Manufacturer               | dianova  |
| Host                       | Goat   |
| Category                   | Antikörper   |
| Application                | DOT, ELISA   |
| Species Reactivity         | Mouse  |
| Immunogen                  | Mouse IgG IgA and IgM whole molecule   |
| Conjugation                | Unconjugated   |
| Product Description        | Anti-Mouse IgG IgA IgM Antibody generated in goat detects reactivity to Mouse IgG, Mouse IgA, and Mouse IgM. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin |
| Clonality                  | Polyclonal   |
| Concentration              | 1.2 mg/mL  |
| Isotype                    | Ig   |

| Buffer             | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2   |
|--------------------|--|
| Purity             | Secondary Antibody was prepared from polyspecific antiserum by immunoaffinity chromatography using antigens coupled to agarose beads followed by cross adsorption to remove unwanted specificities. Assay by immunoelectrophoresis resulted in a single precip |
| Formula            | 20 mM K3PO4,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3   |
| Target             | Mouse  |
| Antibody Type      | Secondary Antibody   |
| Application Dilute | ELISA Dilution: 1:50,000-1:90,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000   |
| Application Notes  | Anti-Mouse IgG IgA IgM antibody has been tested by ELISA and Dot Blot, and is suitable for ELISA, western blot, and immunohistochemistry, as well as other assays requiring lot-to-lot consistency.  |