

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat IgG anti-Mouse IgG (Fc)-unconj., MinX none, Polyclonal DNA-SEC-183159

Article Name Biozol Catalog Number Biozol Catalog Number SEC-183159 Supplier Catalog Number Alternative Catalog Number Manufacturer Manufacturer Most Goat Category Antikörper Application DOT, ELISA, WB Species Reactivity Immunogen Anti-Mouse IgG F(c) fragment was produced by repeated immunization with Mouse IgG F(c) fragment in goat. Conjugation Unconjugated Anti-Mouse IgG F(c) fragment antibody generated in goat detects specifically Mouse IgG F(c) fragment. It is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperatu Clonality Polyclonal Concentration 2.3 mg/mL Isotype Ig		
Supplier Catalog Number Alternative Catalog Number DNA-SEC-183159 Manufacturer dianova Host Goat Category Antikörper Application DOT, ELISA, WB Species Reactivity Mouse Immunogen Anti-Mouse IgG F(c) fragment was produced by repeated immunization with Mouse IgG F(c) fragment in goat. Conjugation Unconjugated Anti-Mouse IgG F(c) fragment antibody generated in goat detects specifically Mouse IgG F(c) fragment. It is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperatu Clonality Polyclonal Concentration 2.3 mg/mL Isotype	Article Name	Goat IgG anti-Mouse IgG (Fc)-unconj., MinX none, Polyclonal
Alternative Catalog Number Manufacturer dianova Host Goat Category Antikörper Application DOT, ELISA, WB Species Reactivity Mouse Anti-Mouse IgG F(c) fragment was produced by repeated immunization with Mouse IgG F(c) fragment in goat. Conjugation Unconjugated Anti-Mouse IgG F(c) fragment antibody generated in goat detects specifically Mouse IgG F(c) fragment. It is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperatu Clonality Polyclonal Concentration 1 January 1981 Lisotype Ig	Biozol Catalog Number	DNA-SEC-183159
Manufacturer Host Goat Category Antikörper Application DOT, ELISA, WB Species Reactivity Mouse Immunogen Anti-Mouse IgG F(c) fragment was produced by repeated immunization with Mouse IgG F(c) fragment in goat. Conjugation Unconjugated Anti-Mouse IgG F(c) fragment antibody generated in goat detects specifically Mouse IgG F(c) fragment. It is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperatu Clonality Polyclonal Concentration 2.3 mg/mL Isotype Ig	Supplier Catalog Number	SEC-183159
Host Goat Category Antikörper Application DOT, ELISA, WB Species Reactivity Mouse Immunogen Anti-Mouse IgG F(c) fragment was produced by repeated immunization with Mouse IgG F(c) fragment in goat. Conjugation Unconjugated Anti-Mouse IgG F(c) fragment antibody generated in goat detects specifically Mouse IgG F(c) fragment. It is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperatu Clonality Polyclonal Concentration 2.3 mg/mL Isotype Ig	Alternative Catalog Number	DNA-SEC-183159
Category Antikörper Application DOT, ELISA, WB Species Reactivity Mouse Immunogen Anti-Mouse IgG F(c) fragment was produced by repeated immunization with Mouse IgG F(c) fragment in goat. Conjugation Unconjugated Anti-Mouse IgG F(c) fragment antibody generated in goat detects specifically Mouse IgG F(c) fragment. It is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperatu Clonality Polyclonal Concentration 2.3 mg/mL Isotype Ig	Manufacturer	dianova
Application DOT, ELISA, WB Species Reactivity Mouse Immunogen Anti-Mouse IgG F(c) fragment was produced by repeated immunization with Mouse IgG F(c) fragment in goat. Conjugation Unconjugated Anti-Mouse IgG F(c) fragment antibody generated in goat detects specifically Mouse IgG F(c) fragment. It is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperatu Clonality Polyclonal Concentration 2.3 mg/mL Isotype Ig	Host	Goat
Species Reactivity Mouse Immunogen Anti-Mouse IgG F(c) fragment was produced by repeated immunization with Mouse IgG F(c) fragment in goat. Conjugation Unconjugated Anti-Mouse IgG F(c) fragment antibody generated in goat detects specifically Mouse IgG F(c) fragment. It is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperatu Clonality Polyclonal Concentration 2.3 mg/mL Isotype Ig	Category	Antikörper
Immunogen Anti-Mouse IgG F(c) fragment was produced by repeated immunization with Mouse IgG F(c) fragment in goat. Conjugation Unconjugated Anti-Mouse IgG F(c) fragment antibody generated in goat detects specifically Mouse IgG F(c) fragment. It is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperatu Clonality Polyclonal Concentration 2.3 mg/mL Isotype Ig	Application	DOT, ELISA, WB
immunization with Mouse IgG F(c) fragment in goat. Conjugation Unconjugated Anti-Mouse IgG F(c) fragment antibody generated in goat detects specifically Mouse IgG F(c) fragment. It is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperatu Clonality Polyclonal Concentration 2.3 mg/mL Isotype Ig	Species Reactivity	Mouse
Anti-Mouse IgG F(c) fragment antibody generated in goat detects specifically Mouse IgG F(c) fragment. It is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperatu Clonality Polyclonal Concentration 2.3 mg/mL Isotype Ig	Immunogen	
Product Description specifically Mouse IgG F(c) fragment. It is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperatu Clonality Polyclonal Concentration 2.3 mg/mL Ig	Conjugation	Unconjugated
Concentration 2.3 mg/mL Isotype Ig	Product Description	specifically Mouse IgG F(c) fragment. It is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the
Isotype Ig	Clonality	Polyclonal
	Concentration	2.3 mg/mL
	Isotype	Ig
Buffer 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2	Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Mouse IgG, Mouse IgG F(c) and Mou
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3
Target	Mouse
Antibody Type	Polyclonal Antibody
Application Dilute	WB: 1:2,000 - 1:10,000
Application Notes	Anti-Mouse IgG F(c) fragment has been tested by ELISA, dot blot, and wester blot and is suitable for use in immunoelectrophoresis, western-blot, competitive western-blot, ELISA and competitive ELISA assays. Specific conditions for reactivity and signal d