

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat IgG anti-Mouse IgG (Fc)-unconj., MinX none, Polyclonal , Unconjugated DNA-SEC-183159

Article Name	Goat IgG anti-Mouse IgG (Fc)-unconj., MinX none, Polyclonal , Unconjugated
Biozol Catalog Number	DNA-SEC-183159
Supplier Catalog Number	SEC-183159
Alternative Catalog Number	DNA-SEC-183159
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	DOT, ELISA, WB
Species Reactivity	Mouse
Immunogen	Anti-Mouse IgG F(c) fragment was produced by repeated immunization with Mouse IgG F(c) fragment in goat.
Conjugation	Unconjugated
Product Description	Anti-Mouse IgG F(c) fragment antibody generated in goat detects specifically Mouse IgG F(c) fragment. It is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperatu
Clonality	Polyclonal
Concentration	2.3 mg/mL
Isotype	Ig

Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Mouse IgG, Mouse IgG F(c) and Mou
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3
Target	Mouse
Antibody Type	Polyclonal Antibody
Application Dilute	ELISA Dilution: 1:20,000 - 1:50,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000
Application Notes	Anti-Mouse IgG F(c) fragment has been tested by ELISA, dot blot, and wester blot and is suitable for use in immunoelectrophoresis, western-blot, competitive western-blot, ELISA and competitive ELISA assays. Specific conditions for reactivity and signal d