

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

**E-Mail:** info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Goat IgG anti-Mouse IgG (Fc)-FITC, MinX none, Polyclonal DNA-SEC-183166

Goat IgG anti-Mouse IgG (Fc)-FITC, MinX none, Polyclonal
DNA-SEC-183166
SEC-183166
DNA-SEC-183166
dianova
Goat
Antikörper
DOT, WB
Mouse
Mouse IgG F(c) fragment
FITC
Anti-Mouse IgG F(c) generated in goat is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc portion of mouse IgG and oft
Polyclonal
2.0 mg/mL
Ig
0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single
Formula	10 mM NaPO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Thimerosal
Target	Mouse
Antibody Type	Polyclonal Antibody
Application Dilute	FLISA 1:10,000 - 1:50,000, FC 1:500 - 1:2,500, IF Microscopy 1:1,000 - 1:5,000
Application Notes	Anti-Mouse IgG F(c) Fluorescein conjugated Antibody has been tested by dot blot and western blot and is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable fo