

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat IgG anti-Mouse IgG (Fc)-Biotin, MinX none, Polyclonal DNA-SEC-183227

| | |
|----------------------------|---|
| Article Name | Goat IgG anti-Mouse IgG (Fc)-Biotin, MinX none, Polyclonal |
| Biozol Catalog Number | DNA-SEC-183227 |
| Supplier Catalog Number | SEC-183227 |
| Alternative Catalog Number | DNA-SEC-183227 |
| Manufacturer | dianova |
| Host | Goat |
| Category | Antikörper |
| Application | WB, IHC, ELISA |
| Species Reactivity | Mouse |
| Immunogen | Anti-mouse IgG F(c) was produced by repeated immunization with mouse IgG F(c) fragment in goat. |
| Conjugation | Biotin |
| Product Description | Anti-Mouse IgG F(c) Antibody generated in goat detects specifically mouse IgG F(c). It is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. R... |
| Clonality | Polyclonal |
| Concentration | 2.0 mg/mL |
| Isotype | Ig |
| Buffer | 0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |

| | |
|--------------------|--|
| Purity | Anti-Mouse IgG F(c) Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resu |
| Formula | 10 mM NaPO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Thimerosal |
| Target | Mouse |
| Antibody Type | Polyclonal Antibody |
| Application Dilute | WB: 1:2,000 - 1:10,000 |
| Application Notes | Anti-Mouse IgG F(c) Antibody has been assayed against 1.0 ug of Mouse IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin S000-03 and ABTS (2,2-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code ABTS-100 as a substrate for 30 |