

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat IgG anti-Rabbit IgG (H+L)-unconj., MinX none, Polyclonal , Unconjugated DNA-SEC-183351

Article Name	Goat IgG anti-Rabbit IgG (H+L)-unconj., MinX none, Polyclonal , Unconjugated
Biozol Catalog Number	DNA-SEC-183351
Supplier Catalog Number	SEC-183351
Alternative Catalog Number	DNA-SEC-183351
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	DOT, ELISA, WB
Species Reactivity	Rabbit
Immunogen	Anti-Rabbit IgG (H&L) was produced by repeated immunization with rabbit whole IgG molecule in goat.
Conjugation	Unconjugated
Product Description	Anti-Rabbit IgG (H&L) Antibody generated in goat detects reactivity to Rabbit IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacte
Clonality	Polyclonal
Concentration	2.20 mg/mL
Isotype	Ig

Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	Rabbit Secondary Antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Rabbit IgG and Rabb
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3
Target	Rabbit
Antibody Type	Polyclonal Antibody
Application Dilute	ELISA Dilution: 1:10,000 - 1:60,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, IF Microscopy Dilution: User Optimized, Immunoprecipitation Dilution: User Optimized, Western Blot Dilution: 1:1,000 - 1:5,000
Application Notes	Anti-Rabbit IgG (H&L) has been tested by ELISA, dot blot, and western blot. This antibody is suitable for use in immunoelectrophoresis, western-blot, competitive western-blot, ELISA and competitive ELISA assays. Specific conditions for reactivity and sig