

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat IgG anti-Rabbit IgG (Fc)-unconj., MinX none, Polyclonal , Unconjugated DNA-SEC-183352

Article Name	Goat IgG anti-Rabbit IgG (Fc)-unconj., MinX none, Polyclonal , Unconjugated
Biozol Catalog Number	DNA-SEC-183352
Supplier Catalog Number	SEC-183352
Alternative Catalog Number	DNA-SEC-183352
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	DOT, ELISA, WB
Species Reactivity	Rabbit
Immunogen	Anti-Rabbit IgG was produced by repeated immunization with rabbit IgG f(c) fragment in goat
Conjugation	Unconjugated
Product Description	Anti-Rabbit IgG F(c) generated in goat is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc portion of rabbit IgG and o
Clonality	Polyclonal
Concentration	2.26 mg/mL
Isotype	Ig

Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Rabbit IgG, Rabbit IgG F(c) and Rabbit
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3
Target	Rabbit
Antibody Type	Polyclonal Antibody
Application Dilute	ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000
Application Notes	Antibody Anti-Rabbit IgG F(c) has been tested by ELISA, dot blot, and western blot and is suitable for immunohistochemistry as well as other peroxidase-antibody based enzymatic assays requiring lot-to-lot consistency.