

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat IgG anti-Rabbit IgG (Fc)-FITC, MinX Hu, Polyclonal DNA-SEC-183362

| Goat IgG anti-Rabbit IgG (Fc)-FITC, MinX Hu, Polyclonal |
|--|
| DNA-SEC-183362 |
| SEC-183362 |
| DNA-SEC-183362 |
| dianova |
| Goat |
| Antikörper |
| FC, IF, FLISA |
| Rabbit |
| Rabbit IgG F(c) fragment |
| FITC |
| Anti-Rabbit IgG F(c) Fluorescein Antibody generated in goat is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc portio |
| Polyclonal |
| 2.0 mg/mL |
| Ig |
| 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
| |

| Purity | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a singl |
|--------------------|--|
| Formula | 20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3 |
| Target | Rabbit |
| Antibody Type | Polyclonal Antibody |
| Application Dilute | FLISA 1:10,000 - 1:50,000, FC 1:500 - 1:2,500, IF Microscopy 1:1,000 - 1:5,000 |
| Application Notes | This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platfor |