

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Goat IgG anti-Rabbit IgG (Fc)-Alk. Phos., MinX Hu, ALP, Polyclonal , AP DNA-SEC-183381

Article Name	Goat IgG anti-Rabbit IgG (Fc)-Alk. Phos., MinX Hu, ALP, Polyclonal , AP
Biozol Catalog Number	DNA-SEC-183381
Supplier Catalog Number	SEC-183381
Alternative Catalog Number	DNA-SEC-183381
Manufacturer	dianova
Host	Goat
Category	Antikörper
Application	DOT, ELISA
Species Reactivity	Rabbit
Immunogen	Rabbit IgG F(c) fragment
Conjugation	AP
Product Description	Anti-Rabbit IgG F(c) alkaline phosphatase antibody generated in goat is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the
Clonality	Polyclonal
Concentration	1.0 mg/mL
Isotype	lg

Buffer	0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a singl
Formula	50 mM TrisHCl,150 mM NaCl,1 mM MgCl,0,1 mM ZnCl,50% (v/v) Glycerol,pH 8,0,sterile filtered,0.05% NaN3
Target	Rabbit
Antibody Type	Polyclonal Antibody
Application Dilute	ELISA Dilution: 1:2,000 - 1:10,000, Immunohistochemistry Dilution: 1:200 - 1:1,000, Western Blot Dilution: 1:500 - 1:2,500
Application Notes	Anti-Rabbit IgG F(c) alkaline phosphatase antibody has been tested by ELISA and dot blot and is assayed against 1.0 $\mu$ g of Rabbit IgG in a standard capture ELISA using pNPP p-nitrophenyl phosphate code NPP-10 as a substrate for 30 minutes at room temper