

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat IgG anti-Rabbit IgG (H+L)-ATTO 647N, MinX Bo,Ck,Go,Gp,Hm,Ho,Hu,Ms,Rt,Sh, ATTO-647N, Polyclonal DNA-SEC-183389

| Article Name | Goat IgG anti-Rabbit IgG (H+L)-ATTO 647N, MinX Bo,Ck,Go,Gp,Hm,Ho,Hu,Ms,Rt,Sh, ATTO-647N, Polyclonal |
|----------------------------|---|
| Biozol Catalog Number | DNA-SEC-183389 |
| Supplier Catalog Number | SEC-183389 |
| Alternative Catalog Number | DNA-SEC-183389 |
| Manufacturer | dianova |
| Host | Goat |
| Category | Antikörper |
| Application | DOT, WB |
| Species Reactivity | Rabbit |
| Immunogen | Rabbit IgG whole molecule |
| Conjugation | ATTO-647N |
| Product Description | Anti-Rabbit IgG (H&L) ATTO 647N Antibody generated in goat detects reactivity to Rabbit IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viru |
| Clonality | Polyclonal |
| Concentration | 1.0 mg/mL |
| Isotype | lg |

| Buffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |
|--------------------|--|
| Purity | Rabbit IgG (H&L) Antibody ATTO 647N was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophore |
| Formula | 20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3 |
| Target | Rabbit |
| Antibody Type | Polyclonal Antibody |
| Application Dilute | FLISA Dilution: >1:20,000, Fluorochrome Protein Value: 1.5, IF Microscopy Dilution: >1:5,000, Western Blot Dilution: >1:10,000 |
| Application Notes | Anti-Rabbit IgG (H&L) conjugated to ATTO 647N has been tested by dot blot and western blot and is designed for STED microscopy, FRET, immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is |