

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Goat IgG anti-Rabbit IgG (H+L)-Biotin, MinX none, Polyclonal DNA-SEC-183393

Goat IgG anti-Rabbit IgG (H+L)-Biotin, MinX none, Polyclonal
DNA-SEC-183393
SEC-183393
DNA-SEC-183393
dianova
Goat
Antikörper
ELISA
Rabbit
Anti-Rabbit IgG whole molecule was produced by repeated immunization with Rabbit IgG whole molecule in goat.
Biotin
Anti-Rabbit IgG (H&L) Biotin Antibody generated in goat detects reactivity to Rabbit IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses
Polyclonal
2.0 mg/mL
Ig
0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2

Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a singl
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3
Target	Rabbit
Antibody Type	Polyclonal Antibody
Application Dilute	ELISA Dilution: 1:50,000 - 1:200,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:5,000 - 1:20,000
Application Notes	Anti-Rabbit IgG Biotin Antibody has been tested by ELISA and is suitable for use in immunoelectrophoresis, IHC, western-blot, competitive western-blot, ELISA and competitive ELISA assays. Specific conditions for reactivity and signal detection should be