

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Donkey IgG anti-Rabbit IgG (H+L)-unconj., MinX Bo,Ck,Go,Gp,Hm,Ho,Hu,Ms,Rt,Sh, Polyclonal, Unconjugated DNA-SEC-183421

Article Name	Donkey IgG anti-Rabbit IgG (H+L)-unconj., MinX Bo,Ck,Go,Gp,Hm,Ho,Hu,Ms,Rt,Sh, Polyclonal , Unconjugated
Biozol Catalog Number	DNA-SEC-183421
Supplier Catalog Number	SEC-183421
Alternative Catalog Number	DNA-SEC-183421
Manufacturer	dianova
Host	Donkey
Category	Antikörper
Application	WB
Species Reactivity	Rabbit
Immunogen	Anti-Rabbit IgG (H&L) was produced by repeated immunization with rabbit whole IgG molecule in donkey.
Conjugation	Unconjugated
Product Description	Anti-Rabbit IgG (H&L) Antibody generated in donkey detects reactivity to Rabbit IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bac
Clonality	Polyclonal
Concentration	1.0 mg/mL
Isotype	Ig

Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a singl
Formula	20 mM K3PO4,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3
Target	Rabbit
Antibody Type	Polyclonal Antibody
Application Dilute	ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000
Application Notes	Anti-Rabbit IgG (H&L) is tested by western blot and suitable for use in immunoelectrophoresis, western-blot, competitive western-blot, ELISA and competitive ELISA assays. Specific conditions for reactivity and signal detection should be optimized by the