

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Donkey IgG anti-Rabbit IgG (Fc)-HRPO, MinX none, Polyclonal DNA-SEC-183424

|                            |   |
|----------------------------|---|
| Article Name               | Donkey IgG anti-Rabbit IgG (Fc)-HRPO, MinX none, Polyclonal   |
| Biozol Catalog Number      | DNA-SEC-183424  |
| Supplier Catalog Number    | SEC-183424  |
| Alternative Catalog Number | DNA-SEC-183424  |
| Manufacturer               | dianova   |
| Host                       | Donkey  |
| Category                   | Antikörper  |
| Application                | ELISA   |
| Species Reactivity         | Rabbit  |
| Immunogen                  | Rabbit IgG F(c) fragment  |
| Conjugation                | HRP   |
| Product Description        | Anti-Rabbit IgG F(c) peroxidase antibody generated in donkey is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme papain under controlled conditions of temperature, time and pH. Receptors bind the Fc porti... |
| Clonality                  | Polyclonal  |
| Concentration              | 2.0 mg/mL   |
| Isotype                    | Ig  |
| Buffer                     | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2  |

|                    |  |
|--------------------|--|
| Purity             | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Donkey Serum, Rabbit IgG, R |
| Formula            | 20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin   |
| Target             | Rabbit   |
| Antibody Type      | Polyclonal Antibody  |
| Application Dilute | ELISA Dilution: 1:10,000 - 1:50,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:1,000 - 1:10,000   |
| Application Notes  | Anti-Rabbit IgG F(c) Peroxidase conjugate has been tested by ELISA and is suitable for immunoblotting (western or dot blot), ELISA, immunoelectron microscopy and immunohistochemistry as well as other antibody-based enzymatic assays requiring lot-to-lot c |