

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Donkey IgG anti-Rabbit IgG (H+L)-Alk. Phos., MinX Bo,Ck,Go,Gp,Hm,Ho,Hu,Ms,Rt,Sh, ALP, Polyclonal, AP DNA-SEC-183427

Article Name	Donkey IgG anti-Rabbit IgG (H+L)-Alk. Phos., MinX Bo,Ck,Go,Gp,Hm,Ho,Hu,Ms,Rt,Sh, ALP, Polyclonal , AP
Biozol Catalog Number	DNA-SEC-183427
Supplier Catalog Number	SEC-183427
Alternative Catalog Number	DNA-SEC-183427
Manufacturer	dianova
Host	Donkey
Category	Antikörper
Application	DOT, ELISA
Species Reactivity	Rabbit
Immunogen	Anti-Rabbit IgG (H&L) was produced by repeated immunization with Rabbit IgG whole molecule in donkey
Conjugation	AP
Product Description	Anti-Rabbit IgG (H&L) Alkaline Phosphatase Antibody generated in donkey detects reactivity to Rabbit IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G
Clonality	Polyclonal
Concentration	1.0 mg/mL
Isotype	Ig

Buffer	0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0
Purity	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a singl
Formula	50 mM TrisHCl,150 mM NaCl,1 mM MgCl,0,1 mM ZnCl,50% (v/v) Glycerol,pH 8,0,sterile filtered,0,01% NaN3
Target	Rabbit
Antibody Type	Polyclonal Antibody
Application Dilute	ELISA Dilution: 1:10,000 - 1:20,000, Immunohistochemistry Dilution: 1:400 - 1:2,000, Western Blot Dilution: 1:1,000 - 1:5,000
Application Notes	Anti-Rabbit IgG (H&L) Alkaline Phosphatase Antibody has been tested by dot blot and ELISA and is suitable for immunoblotting (western or dot blot), ELISA, and immunohistochemistry assays requiring lot-to-lot consistency.