

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat IgG anti-Rat IgG (H+L)-Alk. Phos., MinX Bo,Ck,Go,Gp,Hm,Ho,Hu,Ms,Rb,Sh, ALP, Polyclonal DNA-SEC-183447

| Article Name | Goat IgG anti-Rat IgG (H+L)-Alk. Phos., MinX Bo,Ck,Go,Gp,Hm,Ho,Hu,Ms,Rb,Sh, ALP, Polyclonal |
|----------------------------|---|
| Biozol Catalog Number | DNA-SEC-183447 |
| Supplier Catalog Number | SEC-183447 |
| Alternative Catalog Number | DNA-SEC-183447 |
| Manufacturer | dianova |
| Host | Goat |
| Category | Antikörper |
| Application | DOT, ELISA |
| Species Reactivity | Rat |
| Immunogen | Anti-Rat IgG (H&L) was produced by repeated immunization with Rat IgG whole molecule in goat. |
| Conjugation | ALP |
| Product Description | Anti-Rat IgG (H&L) antibody generated in goat detects specifically rat IgG (H&L). This secondary antibody anti-Rat is ideal for investigators who routinely perform titration assays, western-blot, immunoprecipitation and more generally immunoassays |
| Clonality | Polyclonal |
| Concentration | 1.0 mg/mL |
| Isotype | Ig |

| Buffer | 0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0 |
|--------------------|--|
| Purity | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single p |
| Formula | 50 mM TrisHCl,150 mM NaCl,1 mM MgCl,0,1 mM ZnCl,50% (v/v) Glycerol,pH 8,0,sterile filtered,0,1% NaN3 |
| Target | Rat |
| Antibody Type | Polyclonal Antibody |
| Application Dilute | WB: 1:500 - 1:1,500 |
| Application Notes | Anti-Rat IgG Alkaline Phosphatase has been tested by ELISA and dot blot and is suitable for immunoblotting (western or dot blot), ELISA, and immunohistochemistry assays requiring lot-to-lot consistency. |