

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Goat IgG anti-Rat IgG (H+L)-unconj., MinX Hu, Polyclonal DNA-SEC-183456

|                            |   |
|----------------------------|---|
| Article Name               | Goat IgG anti-Rat IgG (H+L)-unconj., MinX Hu, Polyclonal  |
| Biozol Catalog Number      | DNA-SEC-183456  |
| Supplier Catalog Number    | SEC-183456  |
| Alternative Catalog Number | DNA-SEC-183456  |
| Manufacturer               | dianova   |
| Host                       | Goat  |
| Category                   | Antikörper  |
| Application                | ELISA, WB   |
| Species Reactivity         | Rat   |
| Immunogen                  | Rat IgG whole molecule  |
| Conjugation                | Unconjugated  |
| Product Description        | Anti-Rat IgG (H&L) generated in goat detects rat Immunoglobulin G. Both the Heavy and Light chains of the antibody molecule are present. Representing approximately 75% of serum immunoglobulins, IgG is the most abundant antibody isotype found in the c... |
| Clonality                  | Polyclonal  |
| Concentration              | 2.0 mg/mL   |
| Isotype                    | Ig  |
| Buffer                     | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2  |

|                    |  |
|--------------------|--|
| Purity             | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single p |
| Formula            | 20 mM K3PO4,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3   |
| Target             | Rat  |
| Antibody Type      | Polyclonal Antibody  |
| Application Dilute | WB: 1:2,000 - 1:10,000   |
| Application Notes  | Anti-Rat IgG antibody has been tested by ELISA and western blot and is suitable for use in immunohistochemistry. Specific conditions for reactivity should be optimized by the end user.   |