

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Rabbit IgG anti-Sheep IgG (F(ab)2)-HRPO, MinX Hu, Polyclonal DNA-SEC-183554

|                            |   |
|----------------------------|---|
| Article Name               | Rabbit IgG anti-Sheep IgG (F(ab)2)-HRPO, MinX Hu, Polyclonal  |
| Biozol Catalog Number      | DNA-SEC-183554  |
| Supplier Catalog Number    | SEC-183554  |
| Alternative Catalog Number | DNA-SEC-183554  |
| Manufacturer               | dianova   |
| Host                       | Rabbit  |
| Category                   | Antikörper  |
| Application                | WB, IHC, ELISA  |
| Species Reactivity         | Sheep   |
| Immunogen                  | Sheep IgG F(ab)2 fragment   |
| Conjugation                | HRP   |
| Product Description        | Anti-Sheep IgG F(ab)2 Antibody generated in rabbit recognizes the dimeric Fab portion of the sheep IgG molecule. Sheep IgG F(ab)2 is a proteolytic fragment of immunoglobulin G (IgG) obtained by limited digestion with the enzyme pepsin under controlle... |
| Clonality                  | Polyclonal  |
| Concentration              | 1.0 mg/mL   |
| Isotype                    | Ig  |
| Buffer                     | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2  |

|                    |  |
|--------------------|--|
| Purity             | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Sheep IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single |
| Formula            | 20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin   |
| Target             | Sheep  |
| Antibody Type      | Polyclonal Antibody  |
| Application Dilute | WB: 1:1,000 - 1:5,000  |
| Application Notes  | Anti-Sheep IgG F(ab)2 antibody is suitable for use in ELISA, immunohistochemistry, and western blot. Specific conditions for reactivity should be optimized by the end user.   |