

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat IgG anti-Donkey IgG (H+L)-Biotin, MinX none, Polyclonal DNA-SEC-183597

| | |
|----------------------------|---|
| Article Name | Goat IgG anti-Donkey IgG (H+L)-Biotin, MinX none, Polyclonal |
| Biozol Catalog Number | DNA-SEC-183597 |
| Supplier Catalog Number | SEC-183597 |
| Alternative Catalog Number | DNA-SEC-183597 |
| Manufacturer | dianova |
| Host | Goat |
| Category | Antikörper |
| Application | WB, IHC, ELISA |
| Species Reactivity | Donkey |
| Immunogen | Donkey IgG whole molecule |
| Conjugation | Biotin |
| Product Description | Anti-Donkey IgG Biotin Antibody generated in goat detects donkey IgG. Secreted as part of the adaptive immune response by plasma B cells, immunoglobulin G constitutes 75% of serum immunoglobulins. Immunoglobulin G binds to viruses, bacteria, as well ... |
| Clonality | Polyclonal |
| Concentration | 2.0 mg/mL |
| Isotype | Ig |
| Buffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |

| | |
|--------------------|--|
| Purity | Anti-Donkey IgG (H&L) was prepared from monospecific antiserum by immunoaffinity chromatography using Donkey IgG coupled to agarose. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Goat Serum, Donkey IgG and D |
| Formula | 20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3 |
| Target | Donkey |
| Antibody Type | Polyclonal Antibody |
| Application Dilute | WB: 1:2,000 - 1:10,000 |
| Application Notes | Anti-Donkey IgG (H&L) has been assayed against 1.0 ug of Donkey IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin and ABTS (2,2-azino-bis-[3-ethylbenthiiazoline-6-sulfonic acid]) as a substrate for 30 minutes at room temperature. A |