

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Rabbit IgG anti-Llama IgG2a (Fc)-unconj., MinX none, Polyclonal DNA-SEC-183654

|                            |   |
|----------------------------|---|
| Article Name               | Rabbit IgG anti-Llama IgG2a (Fc)-unconj., MinX none, Polyclonal   |
| Biozol Catalog Number      | DNA-SEC-183654  |
| Supplier Catalog Number    | SEC-183654  |
| Alternative Catalog Number | DNA-SEC-183654  |
| Manufacturer               | dianova   |
| Host                       | Rabbit  |
| Category                   | Antikörper  |
| Application                | DOT, ELISA, WB  |
| Species Reactivity         | Camelus   |
| Immunogen                  | Llama IgG2  |
| Conjugation                | Unconjugated  |
| Product Description        | Camelids produce functional IgG isotypes that do not incorporate light chains. Comparative studies of old world camelids ( <i>Camelus bactrianus</i> and <i>Camelus dromedarius</i> ) and new world camelids ( <i>Lama pacos</i> , <i>Lama glama</i> and <i>Lama vicugna</i> ) have shown that hea... |
| Clonality                  | Polyclonal  |
| Concentration              | 1.07 mg/ml  |
| Isotype                    | Ig  |
| Buffer                     | 0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.2   |

|                    |  |
|--------------------|--|
| Purity             | Anti-Llama IgG2 antibody is directed against Llama IgG2 with minimal cross-reactivity to Llama IgG3 and Llama IgG1. Anti-Llama IgG2 antibody was prepared from monospecific antiserum by immunoaffinity chromatography. Assay by immunoelectrophoresis resulte |
| Formula            | 10 mM NaPO4,250 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3   |
| Target             | Llama  |
| Antibody Type      | Polyclonal Antibody  |
| Application Dilute | WB: 1:20,000-1:40,000  |
| Application Notes  | Antibody Anti-Llama IgG2 has been tested by ELISA, dot blot, and western blot and is suitable for immunoblotting (western or dot blot), ELISA, and immunohistochemistry assays requiring lot-to-lot consistency.   |