

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit F(ab)2 anti-Dog IgG (H+L)-RPE, MinX none, Polyclonal DNA-SEC-183677

| | |
|----------------------------|---|
| Article Name | Rabbit F(ab)2 anti-Dog IgG (H+L)-RPE, MinX none, Polyclonal |
| Biozol Catalog Number | DNA-SEC-183677 |
| Supplier Catalog Number | SEC-183677 |
| Alternative Catalog Number | DNA-SEC-183677 |
| Manufacturer | dianova |
| Host | Rabbit |
| Category | Antikörper |
| Application | FC, IF |
| Species Reactivity | Canine |
| Immunogen | Dog IgG whole molecule |
| Conjugation | RPE |
| Product Description | F(ab)2 Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and exper... |
| Clonality | Polyclonal |
| Concentration | 1.0 mg/mL |
| Isotype | Ig |
| Buffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |

| | |
|--------------------|--|
| Purity | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Dog IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Assa |
| Formula | 20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3 |
| Target | Dog |
| Antibody Type | Polyclonal Antibody |
| Application Dilute | FC 1:100 - 1:250, IF Microscopy 1:100 - 1:250 |
| Application Notes | Secondary antibody reagents are ideal for ELISA, western blotting, Immunohistochemistry, Fluorescence Microscopy, Flow Cytometry as well as other antibody detection methods. |