

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Rabbit F(ab)2 anti-Goat IgG (H+L)-unconj., MinX none, Polyclonal DNA-SEC-183678

| | |
|----------------------------|---|
| Article Name | Rabbit F(ab)2 anti-Goat IgG (H+L)-unconj., MinX none, Polyclonal |
| Biozol Catalog Number | DNA-SEC-183678 |
| Supplier Catalog Number | SEC-183678 |
| Alternative Catalog Number | DNA-SEC-183678 |
| Manufacturer | dianova |
| Host | Rabbit |
| Category | Antikörper |
| Application | WB, IHC, ELISA |
| Species Reactivity | Goat |
| Immunogen | Goat IgG whole molecule |
| Conjugation | Unconjugated |
| Product Description | F(ab)2 Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain immunochemical techniques and exper... |
| Clonality | Polyclonal |
| Concentration | 1.0 mg/mL |
| Isotype | Ig |
| Buffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |

| | |
|--------------------|--|
| Purity | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by pepsin digestion and chromatographic separation. Assay by immunoelectrophoresis resulted in a single precipitin arc |
| Formula | 20 mM K ₃ PO ₄ ,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN ₃ |
| Target | Goat |
| Antibody Type | Polyclonal Antibody |
| Application Dilute | WB: 1:2,000-1:10,000 |
| Application Notes | F(ab) ₂ Anti-Goat IgG Antibody has been tested by SDS-PAGE and is suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated bind |