

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Rabbit F(ab)2 anti-Goat IgG (H+L)-FITC, MinX Hu, Polyclonal DNA-SEC-183685

|                            |   |
|----------------------------|---|
| Article Name               | Rabbit F(ab)2 anti-Goat IgG (H+L)-FITC, MinX Hu, Polyclonal   |
| Biozol Catalog Number      | DNA-SEC-183685  |
| Supplier Catalog Number    | SEC-183685  |
| Alternative Catalog Number | DNA-SEC-183685  |
| Manufacturer               | dianova   |
| Host                       | Rabbit  |
| Category                   | Antikörper  |
| Application                | WB  |
| Species Reactivity         | Goat  |
| Immunogen                  | Goat IgG whole molecule   |
| Conjugation                | FITC  |
| Product Description        | F(ab)2 Anti-Goat IgG Fluorescein Antibody was generated by enzymatic cleavage and subsequent separation from the Fc fragment. Because of their smaller size, F(ab)2 fragments offer several advantages over intact antibodies for use in certain immunoche... |
| Clonality                  | Polyclonal  |
| Concentration              | 1.0 mg/mL   |
| Isotype                    | Ig  |
| Buffer                     | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2  |

|                    |  |
|--------------------|--|
| Purity             | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. Ass |
| Formula            | 20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3   |
| Target             | Goat   |
| Antibody Type      | Polyclonal Antibody  |
| Application Dilute | FLISA 1:10,000 - 1:50,000, FC 1:500-1:2,500, IF Microscopy 1:1,000 - 1:5,000   |
| Application Notes  | F(ab)2 Anti-Goat IgG Fluorescein Antibody has been tested by western blot and is suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F( |