

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat F(ab)2 anti-Human IgG+IgM+IgA (H+L)-Alk. Phos., MinX Ms, ALP, Polyclonal DNA-SEC-183721

| Article Name | Goat F(ab)2 anti-Human IgG+IgM+IgA (H+L)-Alk. Phos., MinX Ms, ALP, Polyclonal |
|----------------------------|--|
| Biozol Catalog Number | DNA-SEC-183721 |
| Supplier Catalog Number | SEC-183721 |
| Alternative Catalog Number | DNA-SEC-183721 |
| Manufacturer | dianova |
| Host | Goat |
| Category | Antikörper |
| Application | WB, IHC, ELISA |
| Species Reactivity | Human |
| Immunogen | Human IgG, IgA, and IgM whole molecules |
| Conjugation | ALP |
| Product Description | F(ab)2 Anti-Human IgG IgA IgM (H&L) Alkaline Phosphatase Antibody generated in goat detects human (heavy and light chain) immunoglobulin G, A, and M. Immunoglobulin G binds to antigens and can neutralize or opsonize targets, and are predominantly inv |
| Clonality | Polyclonal |
| Concentration | 0.5 mg/mL |
| Isotype | Ig |

| Buffer | 0.05 M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol, pH 8.0 |
|--------------------|--|
| Purity | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. As |
| Formula | 50 mM TrisHCl,150 mM NaCl,1 mM MgCl,0,1 mM ZnCl,50% (v/v) Glycerol,pH 8,0,sterile filtered,0,01% NaN3 |
| Target | Human |
| Antibody Type | Polyclonal Antibody |
| Application Dilute | WB: 1:500 - 1:2,500 |
| Application Notes | F(ab)2 Anti-Human IgG IgA IgM (H&L) Alkaline Phosphate Antibody has been assayed against 1.0 ug of Human IgG in a standard capture ELISA using pNPP p-nitrophenyl phosphate code NPP-10 as a substrate for 30 minutes at room temperature. A working dilutio |