

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Goat F(ab)2 anti-Human Kappa light chain-FITC, MinX none, Polyclonal DNA-SEC-183732

|                            |   |
|----------------------------|---|
| Article Name               | Goat F(ab)2 anti-Human Kappa light chain-FITC, MinX none, Polyclonal  |
| Biozol Catalog Number      | DNA-SEC-183732  |
| Supplier Catalog Number    | SEC-183732  |
| Alternative Catalog Number | DNA-SEC-183732  |
| Manufacturer               | dianova   |
| Host                       | Goat  |
| Category                   | Antikörper  |
| Application                | DOT, ELISA, WB  |
| Species Reactivity         | Human   |
| Immunogen                  | Human kappa light chain   |
| Conjugation                | FITC  |
| Product Description        | F(ab)2 HUMAN kappa Antibody Fluorescein Conjugated is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor ima... |
| Clonality                  | Polyclonal  |
| Concentration              | 1.0 mg/ml   |
| Isotype                    | Ig  |
| Buffer                     | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2  |

|                    |  |
|--------------------|--|
| Purity             | F(ab)2 HUMAN kappa Antibody Fluorescein Conjugated was prepared from monospecific antiserum by immunoaffinity chromatography using antigens coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digesti |
| Formula            | 20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3   |
| Target             | Human  |
| Antibody Type      | Polyclonal Antibody  |
| Application Dilute | FLISA 1:10,000 - 1:50,000, FC 1:500 - 1:2,500, IF Microscopy 1:1,000 - 1:5,000   |
| Application Notes  | F(ab)2 Human kappa Fluorescein Conjugated Antibody has been tested by ELISA, dot blot, and western blot and is suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low ba |