

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat F(ab)2 anti-Human IgG (Fc)-HRPO, MinX Bo,Ho,Ms,Rt, Polyclonal DNA-SEC-183742

| | |
|----------------------------|---|
| Article Name | Goat F(ab)2 anti-Human IgG (Fc)-HRPO, MinX Bo,Ho,Ms,Rt, Polyclonal |
| Biozol Catalog Number | DNA-SEC-183742 |
| Supplier Catalog Number | SEC-183742 |
| Alternative Catalog Number | DNA-SEC-183742 |
| Manufacturer | dianova |
| Host | Goat |
| Category | Antikörper |
| Application | ELISA |
| Species Reactivity | Human |
| Immunogen | Human IgG F(c) fragment |
| Conjugation | HRP |
| Product Description | F(ab)2 Anti-Human IgG F(c) Peroxidase Antibody generated in goat detects Human F(c). Representing approximately 75% of serum immunoglobulins in humans, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesized ... |
| Clonality | Polyclonal |
| Concentration | 1.0 mg/mL |
| Isotype | Ig |
| Buffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |

| | |
|--------------------|--|
| Purity | F(ab)2 Anti-Human IgG Fc was prepared from monospecific antiserum by immunoaffinity chromatography using Human IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic se |
| Formula | 20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin |
| Target | Human |
| Antibody Type | Polyclonal Antibody |
| Application Dilute | WB: 1:1,000 - 1:5,000 |
| Application Notes | F(ab)2 Anti-Human IgG F(c) Peroxidase Antibody has been tested by ELISA and is suitable for use in immunoelectrophoresis, western-blot, competitive western-blot, ELISA and competitive ELISA assays. Specific conditions for reactivity and signal detection |