

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### Goat F(ab)2 anti-Mouse IgG (F(ab)2)-unconj., MinX Bo,Ho,Hu,Rb,Rt,Sh, Polyclonal DNA-SEC-183782

|                            |   |
|----------------------------|---|
| Article Name               | Goat F(ab)2 anti-Mouse IgG (F(ab)2)-unconj., MinX Bo,Ho,Hu,Rb,Rt,Sh, Polyclonal   |
| Biozol Catalog Number      | DNA-SEC-183782  |
| Supplier Catalog Number    | SEC-183782  |
| Alternative Catalog Number | DNA-SEC-183782  |
| Manufacturer               | dianova   |
| Host                       | Goat  |
| Category                   | Antikörper  |
| Application                | ELISA   |
| Species Reactivity         | Mouse   |
| Immunogen                  | Mouse IgG F(ab)2 fragment   |
| Conjugation                | Unconjugated  |
| Product Description        | F(ab)2 Anti-Mouse IgG F(ab)2 Antibody generated in goat detects Mouse F(ab)2. Representing approximately 75% of serum immunoglobulins, IgG is the most abundant antibody isotype found in the circulation. IgG molecules are synthesized and secreted by p... |
| Clonality                  | Polyclonal  |
| Concentration              | 1.0 mg/mL   |
| Isotype                    | Ig  |

|                    |  |
|--------------------|--|
| Buffer             | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2   |
| Purity             | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, pepsin digestion and chromatographic separation. As |
| Formula            | 20 mM K3PO4,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3   |
| Target             | Mouse  |
| Antibody Type      | Polyclonal Antibody  |
| Application Dilute | WB: 1:2,000 - 1:10,000   |
| Application Notes  | F(ab)2 Anti-Mouse IgG F(ab)2 Antibody has been tested by ELISA and is suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated |