

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Goat F(ab)2 anti-Rabbit IgG (F(ab)2)-HRPO, MinX Bo,Ho,Hu,Ms,Rt,Sh, Polyclonal DNA-SEC-183834

| Article Name               | Goat F(ab)2 anti-Rabbit IgG (F(ab)2)-HRPO, MinX Bo,Ho,Hu,Ms,Rt,Sh,<br>Polyclonal   |
|----------------------------|--|
| Biozol Catalog Number      | DNA-SEC-183834   |
| Supplier Catalog Number    | SEC-183834   |
| Alternative Catalog Number | DNA-SEC-183834   |
| Manufacturer               | dianova  |
| Host                       | Goat   |
| Category                   | Antikörper   |
| Application                | ELISA, WB  |
| Species Reactivity         | Rabbit   |
| Immunogen                  | Anti-Rabbit IgG F(ab)2 Antibody was produced in goat by repeated immunization with Rabbit IgG F(ab)2 fragment.   |
| Conjugation                | HRP  |
| Product Description        | F(ab)2 Anti-Rabbit IgG F(ab)2 Peroxidase Antibody generated in goat<br>detects Rabbit F(ab)2. Representing approximately 75% of serum<br>immunoglobulins, IgG is the most abundant antibody isotype found<br>in the circulation. IgG molecules are synthesized and |
| Clonality                  | Polyclonal   |
| Concentration              | 1.0 mg/mL  |
| Isotype                    | lg   |

| Buffer             | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2  |
|--------------------|---|
| Purity             | Anti-Rabbit IgG F(ab)2 Antibody was prepared from monospecific<br>antiserum by immunoaffinity chromatography using Rabbit IgG<br>coupled to agarose beads followed by solid phase adsorption(s) to<br>remove any unwanted reactivities, pepsin digestion and chromatogr |
| Formula            | 20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin  |
| Target             | Rabbit  |
| Antibody Type      | Polyclonal Antibody   |
| Application Dilute | WB: 1:2,000 - 1:10,000  |
| Application Notes  | F(ab)2 Anti-Rabbit IgG F(ab)2 Peroxidase Antibody has been tested<br>by ELISA and western blot and is suitable for immunoblotting<br>(western or dot blot), ELISA, immunoperoxidase electron microscopy<br>and immunohistochemistry as well as other peroxidase-antibod |