

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

**Telephone:** +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Rabbit Fab anti-Goat IgG (H+L)-HRPO, MinX none, Polyclonal DNA-SEC-183925

| Article Name               | Rabbit Fab anti-Goat IgG (H+L)-HRPO, MinX none, Polyclonal   |
|----------------------------|--|
| Biozol Catalog Number      | DNA-SEC-183925   |
| Supplier Catalog Number    | SEC-183925   |
| Alternative Catalog Number | DNA-SEC-183925   |
| Manufacturer               | dianova  |
| Host                       | Rabbit   |
| Category                   | Antikörper   |
| Application                | WB, IHC, ELISA   |
| Species Reactivity         | Goat   |
| Immunogen                  | Goat IgG whole molecule  |
| Conjugation                | HRP  |
| Product Description        | Fab Anti-Goat IgG Peroxidase Antibody generated in rabbit detects goat IgG. This product possesses the F(ab) region possessing the epitope-recognition site, both heavy and light chains of the antibody molecule are present. Secondary Antibodies are av |
| Clonality                  | Polyclonal   |
| Concentration              | 1.0 mg/mL  |
| Isotype                    | Ig   |
| Buffer                     | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2   |

| Purity             | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, papain digestion and chromatographic separation. Ass |
|--------------------|--|
| Formula            | 20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin   |
| Target             | Goat   |
| Antibody Type      | Polyclonal Antibody  |
| Application Dilute | WB: 1:1,000 - 1:4,000  |
| Application Notes  | Suitable for immunoblotting (western or dot blot), ELISA, immunoperoxidase electron microscopy and immunohistochemistry as well as other peroxidase-antibody based enzymatic assays requiring extremely low background levels, absence of F(c) mediated bindin |