

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | Fax: +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## **Product Datasheet**

## Goat Fab anti-Hamster generally IgG (H+L)-Biotin, MinX none, Polyclonal DNA-SEC-183942

| Article Name               | Goat Fab anti-Hamster generally IgG (H+L)-Biotin, MinX none,<br>Polyclonal   |
|----------------------------|--|
| Biozol Catalog Number      | DNA-SEC-183942   |
| Supplier Catalog Number    | SEC-183942   |
| Alternative Catalog Number | DNA-SEC-183942   |
| Manufacturer               | dianova  |
| Host                       | Goat   |
| Category                   | Antikörper   |
| Application                | WB, IHC, ELISA   |
| Species Reactivity         | Golden Hamster   |
| Immunogen                  | Golden Syrian Hamster IgG whole molecule   |
| Conjugation                | Biotin   |
| Product Description        | Fab Anti-Golden Syrian Hamster IgG (H&L) Antibody generated in goat detects immunoglobulin g from hamster, both heavy and light chains of the antibody molecule are present. Each IgG has two antigen binding sites. Representing approximately 75% of ser |
| Clonality                  | Polyclonal   |
| Concentration              | 1.0 mg/mL  |
| Isotype                    | Ig   |
| Buffer                     | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2   |

| Purity             | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Golden Syrian Hamster IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, papain digestion and chromatographi |
|--------------------|--|
| Formula            | 20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% NaN3   |
| Target             | Golden Syrian Hamster  |
| Antibody Type      | Polyclonal Antibody  |
| Application Dilute | WB: 1:2,000 - 1:10,000   |
| Application Notes  | Suitable for immunoblotting, ELISA, immunohistochemistry, immunomicroscopy as well as other antibody based assays using streptavidin or avidin conjugates requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, |