

Diagnostica Vertrieb GmbH, Leipziger Straße 4

85386 Eching, Germany

Telephone: +49 (0)89 3799666-6 | **Fax:** +49 (0)89 3799666-99

E-Mail: info@biozol.de

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

Product Datasheet

Goat Fab anti-Rat IgG (H+L)-HRPO, MinX none, Polyclonal DNA-SEC-183974

| Article Name | Goat Fab anti-Rat IgG (H+L)-HRPO, MinX none, Polyclonal |
|----------------------------|--|
| Biozol Catalog Number | DNA-SEC-183974 |
| Supplier Catalog Number | SEC-183974 |
| Alternative Catalog Number | DNA-SEC-183974 |
| Manufacturer | dianova |
| Host | Goat |
| Category | Antikörper |
| Application | ELISA |
| Species Reactivity | Rat |
| Immunogen | Anti-Rat IgG (H&L) was produced by repeated immunization with rat IgG whole molecule in goat. |
| Conjugation | HRP |
| Product Description | Fab Anti-Rat IgG HRP conjugated secondary antibody generated in goat detects specifically rat IgG. This secondary peroxidase conjugated antibody anti-Rat is ideal for investigators who routinely perform titration assays, western blot, immunoprecipita |
| Clonality | Polyclonal |
| Concentration | 1.0 mg/mL |
| Isotype | Ig |
| Buffer | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 |

| Purity | Anti-Rat IgG (H&L) peroxidase conjugated antibody was prepared from monospecific antiserum by immunoaffinity chromatography using Rat IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, papain digestion |
|--------------------|--|
| Formula | 20 mM K3PO4,150 mM NaCl,pH 7,2,lyophilisate,0,01% Gentamicin |
| Target | Rat |
| Antibody Type | Polyclonal Antibody |
| Application Dilute | ELISA Dilution: 1:10,000 - 1:40,000, Immunohistochemistry Dilution: 1:500 - 1:2,500, Western Blot Dilution: 1:1,000 - 1:4,000 |
| Application Notes | Anti-Rat IgG HRP antibody has been tested by ELISA and is suitable for immunoblotting (western or dot blot), ELISA, immunoperoxidase electron microscopy and immunohistochemistry as well as other peroxidase-antibody based enzymatic assays requiring extrem |