

Please note: This document was created automatically and is not a substitute for the manufacturer's original document.

## Product Datasheet

### **Rabbit Fab anti-Sheep IgG (H+L)-unconj., MinX none, Polyclonal , Unconjugated DNA-SEC-183980**

|                            |   |
|----------------------------|---|
| Article Name               | Rabbit Fab anti-Sheep IgG (H+L)-unconj., MinX none, Polyclonal ,<br>Unconjugated  |
| Biozol Catalog Number      | DNA-SEC-183980  |
| Supplier Catalog Number    | SEC-183980  |
| Alternative Catalog Number | DNA-SEC-183980  |
| Manufacturer               | dianova   |
| Host                       | Rabbit  |
| Category                   | Antikörper  |
| Application                | WB, IHC, ELISA  |
| Species Reactivity         | Sheep   |
| Immunogen                  | Sheep IgG whole molecule  |
| Conjugation                | Unconjugated  |
| Product Description        | Fab Anti-Sheep IgG (H&L) Antibody generated in rabbit detects immunoglobulin g from sheep, both heavy and light chains of the antibody molecule are present. Each IgG has two antigen binding sites. Representing approximately 75% of serum immunoglobuli... |
| Clonality                  | Polyclonal  |
| Concentration              | 0.5 mg/mL   |
| Isotype                    | Ig  |

|                    |  |
|--------------------|--|
| Buffer             | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2   |
| Purity             | This product was prepared from monospecific antiserum by immunoaffinity chromatography using Sheep IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities, papain digestion and chromatographic separation. As |
| Formula            | 20 mM K3PO4,150 mM NaCl,pH 7,2,sterile filtered,0,01% NaN3   |
| Target             | Sheep  |
| Antibody Type      | Polyclonal Antibody  |
| Application Dilute | ELISA Dilution: 1:20,000 - 1:100,000, Immunohistochemistry Dilution: 1:1,000 - 1:5,000, Western Blot Dilution: 1:2,000 - 1:10,000  |
| Application Notes  | Suitable for highly specific immunological methods requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.  |